

New records of six *Pyrenula* species from China

JIE-MENG FU ^{1#}, ZHONG-LIANG WANG ^{1#},
CHUN-XIAO WANG ¹, LU-LU ZHANG ^{2*}

¹Key Laboratory of Plant Stress Research, College of Life Sciences,
Shandong Normal University, Jinan, 250014, P. R. China

²Institute of Environment and Ecology, Shandong Normal University,
Jinan, 250014, P. R. China

* CORRESPONDENCE TO: jiemengfu11@outlook.com, wzl010118@outlook.com

ABSTRACT—*Pyrenula mastophora*, *P. micheneri*, and *P. thailandica* are reported for the first time from China; and *P. bahiana*, *P. complanata*, and *P. platystoma* are reported for the first time from mainland China. Descriptions, illustrations, and distributions are given for each species.

KEY WORDS—Asia, lichen-forming fungi, *Pyrenulaceae*, taxonomy

Introduction

The conserved genus *Pyrenula* Ach. (*Pyrenulaceae*, *Ascomycota*) was described by Acharius (1814). In a world key of *Pyrenula* species, Aptroot (2012) accepted 169 species out of the c. 745 named taxa in the genus. Since then, many new species of *Pyrenula* have been described and the genus now comprises c. 226 species (Ingle & al. 2018).

Pyrenula comprises crustose lichens typically growing on smooth, shaded bark (Aptroot 2012). It is characterized by a UV– or UV+ yellow thallus, perithecioid ascomata, an occasionally interspersed hamathecium, unbranched filaments, and distoseptate or (sub)muriform brown mature ascospores, and old spores with or without orange oil granules. Aptroot & al. (2013) noted that

[#] Jie-Meng Fu & Zhong-Liang Wang contributed equally to this paper.

the presence of orange reddish granules in over-mature spores is an important character that has made it much easier to recognize undescribed taxa.

In this paper we contribute to the knowledge of *Pyrenula* in China by providing fundamental data and reliable results in preparation for the Lichen Flora of China. We have identified three species new to China—*Pyrenula mastophora*, *P. micheneri*, and *P. thailandica*—and report for the first time three species from mainland China—*P. bahiana*, *P. complanata*, and *P. platystoma*.

Materials & methods

The specimens studied are preserved in Lichen Section of Botanical Herbarium, Shandong Normal University, Jinan, China (SDNU). The morphological and anatomical characters of the specimens were examined under a COIC XTL7045B2 stereo-microscope and an Olympus CX41 polarizing microscope. Unless otherwise indicated, sections were mounted in tap water (unless otherwise indicated); measurements were made in tap water or water enhanced with KOH. Lichen substances were identified using standardized thin layer chromatography (TLC) with system C (Orange & al. 2010). The lichens were photographed using Olympus SZX16 and BX61 microscopes and an Olympus DP72 camera.

Taxonomy

Pyrenula bahiana Malme, Ark. Bot. 22A(11): 26 (1929)

FIG. 1A–D

Thallus corticate, brownish to olive green, with pseudocyphellae, thallus UV–. Perithecia solitary, subglobose, immersed to erumpent from the substratum, often partly or mostly covered by thallus, 0.5–0.7 mm diam. Perithecial wall with crystals, up to 100 µm thick, often not fully carbonized. Ostiole pale, apical. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 8/ascus, grey brown, irregularly biseriate, 3-septate, fusiform with pointed ends, $27.5\text{--}30 \times 10\text{--}15$ µm, lumina angular to usually somewhat rounded, all lumina separated from the exospore by a thick endospore layer, old ascospores filled with red-orange oil. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMENS EXAMINED: CHINA. GUIZHOU, Jiangkou, Mt. Xiaoding, alt. 900 m, on bark, 4 Apr. 2016, Xiang-Xiang Zhao 20160694 (SDNU); alt. 850 m, on bark, 2 Apr. 2016, Xiang-Xiang Zhao 2016080 (SDNU).

DISTRIBUTION—Pantropical (Aptroot 2012). New to mainland China, but previously reported from Taiwan (Aptroot 2003).

COMMENTS—*Pyrenula bahiana* is characterized by the thallus with white pseudocyphellae, distoseptate ascospores with three septa, and over-mature




FIG. 1 *Pyrenula bahiana* (Zhao 20160694, SDNU). A: thallus with ascomata and pseudocyphellae; B: transverse section through ascoma; C: ascospores; D: over-mature spores with red-orange oil. *Pyrenula complanata* (Wang 20107042, SDNU). E: thallus with ascomata; F: transverse section through ascoma; G: ascus; H: ascospores. *Pyrenula mastophora* (Li 20110618, SDNU). I: thallus with ascomata; J: transverse section through ascoma; K: ascospores; L: ascus with 8 ascospores.

spores containing red-orange oil. It is closely related to *P. thailandica*, which has larger ascospores ($37.5\text{--}42.5 \times 15\text{--}17.5 \mu\text{m}$). Our specimens are similar to the *P. bahiana* protologue description, except that the holotype has larger ($29\text{--}38 \times 13\text{--}18 \mu\text{m}$) ascospores (Ingle & al. 2018).

Pyrenula complanata (Mont.) Trevis., Spighe e Paglie: 17 (1853) FIG. 1E–H

Thallus corticate, pale brownish, without pseudocyphellae, thallus UV-. Perithecia mostly simple, only aggregated as by chance when crowded, erumpent from the substratum, perithecia subglobose and spreading laterally, 0.9–1.4 mm diam. Perithecial wall <225 µm. Ostiole black, apical. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 8/ascus, grey brown, 3-septate, ellipsoid with rounded ends, 30–32.5 × 10–12.5 µm, end lumina separated from the exospore by a thick endospore layer, lumina angular. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMENS EXAMINED: CHINA. GUANGXI, Laibinjinxiu, Mt. Shengtang, alt. 1400 m, on bark, 8 Nov. 2010, Hai-Ying Wang 20107042 (SDNU). FUJIAN, Longyan, Mt. Huanglianyu, alt. 600 m, on bark, 2 Apr. 2010, Xing-Ran Kou 20105891 (SDNU). YUNNAN, Jingdong, Mt. Ailao, Dujuanhu, alt. 2500 m, on bark, 5 Aug. 2017, Rong Tang & al. 20170335 (SDNU).

DISTRIBUTION—Pantropical (Aptroot 2012). New to mainland China, but previously reported from Taiwan (Aptroot 2003).

COMMENTS—*Pyrenula complanata* is characterized by the thallus lacking pseudocyphellae and distoseptate ascospores with three septa. It is closely related to *P. media*, which has shorter ascospores (24–29 µm) and a thicker thallus (Aptroot 2012). Our specimens are closely similar to the *P. complanata* protologue description (Montagne 1843).

Pyrenula mastophora (Nyl.) Müll. Arg., Flora 66: 426 (1883) FIG. 1I–L

Thallus corticate, pale brownish, without pseudocyphellae, thallus UV-. Perithecia simple, solitary, erumpent from the substratum, 0.7–0.9 mm diam. Perithecial wall up to 200 µm. Ostiole black, apical. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 8/ascus, grey brown, 3-septate, ellipsoid with rounded ends, 25–27.5 × 10–12.5 µm, Central lumina much wider than long, end lumina separated from the exospore by a thick endospore layer. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMENS EXAMINED: CHINA. GUANGXI, Baise, Tianlin, Mt. Cenwanglao, alt. 2000 m, on bark, 23 Mar. 2011, Lin Li 20110618 (SDNU). GUIZHOU, Kaili, Mt. Leigong, alt. 2130 m, on bark, 1 Nov. 2009, Zun-Tian Zhao (SDNU).

DISTRIBUTION—Pantropical (Aptroot 2012). New to China.

COMMENTS—*Pyrenula mastophora* is characterized by the thallus without pseudocyphellae, distoseptate ascospores with three septa and wide central




FIG. 2 *Pyrenula micheneri* (Li 20106097, SDNU). A: thallus with ascomata; B: transverse section through ascoma; C: ascus; D: ascospores. *Pyrenula platystoma* (Tang & al. 20171074, SDNU). E: thallus with ascomata; F: transverse section through ascoma; G: Immature ascospore; H: mature ascospore. *Pyrenula thailandica* (Wang 20102341, SDNU). I: thallus with ascomata and pseudocyphellae; J: ascospores; K: over-mature spores with red-orange oil.

lumina, and non-conical ascomata. It is closely related to *P. quassiicola*, which has more or less rounded central lumina and often possesses pseudocyphellae. Our specimens match the *P. mastophora* protologue description, except that the holotype has conical ascomata (Aptroot 2012).

Pyrenula micheneri R.C. Harris,

Mem. New York Bot. Gard. 49: 96 (1989)

FIG. 2A–D

Thallus corticate, pale brownish to olive green, without pseudocyphellae, thallus UV–. Perithecia simple, only aggregated as by chance when crowded, immersed to erumpent from the substratum, often partly or mostly covered by thallus, perithecia subglobose, 0.4–0.6 mm diam. Ostiole pale, apical, with numerous crystals around ostioles. Perithecial wall with crystals, <150 µm. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 8/ascus, grey brown, 3-septate, ellipsoid with rounded ends, with angular diamond-shaped lumina, 22.5–27.5 × 7.5–10 µm, end lumina separated from the exospore by a thick endospore layer. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMEN EXAMINED: CHINA. JIANGXI, Jian, Qianmo, Nanfengmian, alt. 1300 m, on bark, 2 Nov. 2010, Ming Li 20106097 (SDNU).

DISTRIBUTION—Temperate America (Aptroot 2012). New to China.

COMMENTS—*Pyrenula micheneri* is characterized by a thallus without pseudocyphellae, an apical ostiole, numerous crystals surrounding the ostiole, and distoseptate ascospores with three septa and angular diamond-shaped lumina. It is closely related to *P. pyrgillospora*, which has rounded or quadrangular lumina (Aptroot 2012). Our specimen closely matches the *P. micheneri* protologue description (Harris 1989).

Pyrenula platystoma (Müll. Arg.) Aptroot, Lichenologist 44: 36 (2012) FIG. 2E–H

Thallus corticate, yellowish brown, without pseudocyphellae, thallus UV–. Perithecia simple, only aggregated as by chance when crowded, erumpent from the substratum, often partly covered by thallus, 0.7–1.0 mm diam. Perithecial wall <250 µm. Ostiole pale, apical. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 2/ascus, grey brown, muriform with 8–10 rows, fusiform with rounded ends, 145–150 × 28.5–30 µm. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMEN EXAMINED: CHINA. YUNNAN, Midu, Mt. Taijiding, alt. 2650 m, on bark, 8 Aug. 2017, Rong Tang & al. 20171074 (SDNU).

DISTRIBUTION—Pantropical (Aptroot 2012). New to mainland China, but previously reported from Taiwan as *Anthracothecium oculatum* Müll. Arg. (Zahlbruckner 1940, Lamb 1963, Wang & Lai 1973, Kashiwadani & Kurokawa 1981, Wei 1991), which is a synonym of *P. platystoma* (Aptroot 2012).

COMMENTS—*Pyrenula platystoma* is characterized by the thallus without pseudocyphellae and large muriform ascospores. It is closely related to *P. neosandwicensis*, which has pseudocyphellae (Aptroot 2012). Our specimen matches the *P. platystoma* protologue description, except that the holotype has mostly shorter (80–140(–155) μm long) ascospores (Aptroot 2012).

Pyrenula thailandica Aptroot, Lichenologist 44: 617 (2012)

FIG. 2I–K

Thallus corticate, brownish green, with pseudocyphellae, thallus UV–. Perithecia solitary, subglobose, erumpent from the substratum, 0.7–0.8 mm diam. Ostiole pale, apical. Hamathecium not interspersed with oil droplets, filaments unbranched. Ascospores 8/ascus, grey brown, 3-septate, fusiform with pointed ends, $37.5\text{--}42.5 \times 15\text{--}17.5 \mu\text{m}$, lumina angular to usually somewhat rounded, end lumina separated from the exospore by a thick endospore layer, over-mature ascospores filled with red-orange oil. Substrate bark.

CHEMISTRY—No chemical substances detected by TLC.

SPECIMEN EXAMINED: CHINA. GUIZHOU, Libo, Maolan protection zone, alt. 500 m, on bark, 2 Nov. 2009, Hai-Ying Wang 20102341 (SDNU).

DISTRIBUTION—Eastern palaeotropical (Aptroot & al. 2012). New to China.

COMMENTS—*Pyrenula thailandica* is characterized by the thallus with white pseudocyphellae, distoseptate ascospores with three septa, and old spores with red-orange oil. It is closely related to *P. bahiana*, which has smaller ascospores ($25\text{--}27.5 \times 10\text{--}15 \mu\text{m}$). Our specimen matches the *P. thailandica* protologue description, except that the holotype has $(30\text{--})35\text{--}51 \times (10\text{--})14\text{--}20 \mu\text{m}$ ascospores (Aptroot & al. 2012).

Acknowledgements

The authors thank Dr. Aptroot A (ABL Herbarium, The Netherlands) and Dr. Shou-Yu Guo (State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing) for specimen identification guidance and presubmission reviews. We also thank Dr. Aptroot for providing great help during the study. This study was supported by the National Natural Science Foundation of China (31400015, 31570017) and the Undergraduate Scientific Research Foundation of Shandong Normal University (2017BKSKY61).

Literature cited

- Acharius, E. 1814. *Synopsis methodica lichenum*. Lund. 392 p.
Aptroot A. 2003. Pyrenocarpous lichens and related non-lichenized ascomycetes from Taiwan. *Journal of the Hattori Botanical Laboratory* 93: 155–173.
Aptroot A. 2012. A world key to the species of *Anthracotheceum* and *Pyrenula*. *Lichenologist* 44: 5–53. <https://doi.org/10.1017/S0024282911000624>

- Aptroot A, Schumm F, Cáceres MES. 2012. Six new species of *Pyrenula* from the tropics. *Lichenologist* 44: 611–618. <https://doi.org/10.1017/S0024282912000254>
- Aptroot A, Sipman HJM, Cáceres MES. 2013. Twenty-one new species of *Pyrenula* from South America, with a note on over-mature ascospores. *Lichenologist* 45: 169–198. <https://doi.org/10.1017/S0024282912000734>
- Harris R.C. 1989. A sketch of the family *Pyrenulaceae* (*Melanommatales*) in eastern North America. *Memoirs of the New York Botanical Garden* 49: 74–107.
- Ingle KK, Uppadhyay V, Nayaka S, Trivedi S, Sahoo D. 2018. New records and an updated key of *Pyrenula* from India. *Cryptogam Biodiversity and Assessment*. 1: 37–46. <https://doi.org/10.21756/cab.esp7>
- Kashiwadani H, Kurokawa S. 1981. Notes on Japanese and Formosan species of *Anthracothecium* (2). *Journal of Japanese Botany* 56(11): 348–356.
- Lamb IM. 1963. *Index Nominum Lichenum inter annos 1932 et 1960 divulgatorum*. New York.
- Montagne JPFC. 1843. Troisième centurie de plantes cellulaires exotiques nouvelles. *Décade IX et X. Annales des Sciences Naturelles, Botanique* 19: 53–85.
- Orange A, James PW, White FJ. 2010. *Microchemical methods for the identification of lichens*. 2nd edition. British Lichen Society, London.
- Wang-Yang JR, Lai MJ. 1973. A checklist of the lichens of Taiwan. *Taiwania* 18(1): 83–104.
- Wei JC. 1991. *An enumeration of lichens in China*. International Academic Publishers, Beijing.
- Zahlbruckner A. 1940. *Catalogus Lichenum Universalis*. 1. Leipzig. (Reprinted by Johnson Reprint Corporation, New York).